

BronchoscopyDischarge Instructions

You have received medicine to help you relax during the test.

For your own and other's safety please follow these guidelines:

- A responsible adult MUST accompany you home after your procedure (you cannot go home alone or unaccompanied in a taxi).
- We recommend you have someone stay with you for the first 24 hours that you are home.
- For 24 hours after an anaesthetic, (until _____ am/pm tomorrow) do not do anything that needs coordination or judgment. This includes:
 - Driving a car or operating dangerous machinery.
 - Making important decisions.
 - Signing legal papers.
- Do not drink alcohol or take recreational drugs for 24 hours afterwards. These substances may add to the anaesthetic effects.

Resume fluids:	

When tolerating fluids well, have a light diet for today, resume you regular diet tomorrow.

Please restart your usual prescribed medications immediately unless your Doctor gives you different instructions.

Do not take herbal or Chinese medicines for 1 week after your procedure.

...see over

If you have any of the following:

- Pain in the chest or problems swallowing. (You may have a slight sore throat.)
- · Shortness of breath or pain when you breathe.
- Coughing up blood. (You may have a slight amount of blood in your sputum for a short time after your test. This is normal.)
- Fever (T > 38.5° C) or chills. These may happen the evening of the test or the next day.

CALL THE DOCTOR WHO DID YOUR TEST IMMEDIATELY AT 604-875-4111.

** If your doctor is not available, ask to speak to the Respirologist on call.

If you cannot reach either doctor, go to the closest Emergency Department.**

HealthLink BC

To speak with a nurse, pharmacist, or dietician, call 811. Services available in 130 languages.

www.healthlink.bc.ca

For deaf and hearing-impaired assistance (TTY), call 711.